

A REMARK ON THE TRIGONOMETRIC SYSTEM

ALEXANDER KUSHPEL

ABSTRACT. We present a new property of the trigonometric system arranged in a natural order. It is shown that the sequence of subspaces of trigonometric polynomials $\mathcal{T}_n$ is optimal in the sense of order of convergence on convolution classes $K * U_p$ in L_q for any $1 < p, q < \infty$ just in the cases of "very slow" or "very fast" rate of decay of Fourier coefficients of K .

1. INTRODUCTION

Let X be a Banach space and A a convex, compact, centrally symmetric subset of X . The Kolmogorov n -width of A in X is defined by

$$d_n(A, X) := \inf_{X_n \subset X} \sup_{f \in A} \inf_{g \in X_n} \|f - g\|_X,$$

where X_n runs over all subspaces of X of dimension $\leq n$. Let $\mathbb{T}^1$ be the unit circle with the Lebesgue measure dx . The space of p -integrable functions ϕ , i.e. such that $\|\phi\|_p < \infty$ is denoted by $L_p(\mathbb{T}^1) \equiv L_p$, where

$$\|\phi\|_p := \left(\int_{\mathbb{T}^1} |\phi(x)|^p dx \right)^{1/p}, \quad 1 \leq p < \infty.$$

Consider the sequence of subspaces $\mathcal{T}_n$ of trigonometric polynomials with the usual order,

$$\mathcal{T}_0 := \text{lin}\{1\}, \dots, \mathcal{T}_n := \text{lin}\{1, \cos kx, \sin kx, 1 \leq k \leq n\}, \dots$$

Let $\phi \in L_p$ with the formal Fourier series

$$\phi \sim \sum_{k=1}^{\infty} a_k(\phi) \cos kx + b_k(\phi) \sin kx.$$

A wide range of sets of smooth functions on $\mathbb{T}^1$ can be introduced using multipliers $\Lambda := \{\lambda_k, k \in \mathbb{N}\}$ (see, e.g. [6]). We say that $f \in \Lambda_\beta U_p$, $\beta \in \mathbb{R}$ if

$$f \sim \sum_{k=1}^{\infty} \lambda_k \left(a_k(\phi) \cos \left(kx - \frac{\beta\pi}{2} \right) + b_k(\phi) \sin \left(kx - \frac{\beta\pi}{2} \right) \right),$$

where $\phi \in U_p := \left\{ \phi \mid \|\phi\|_p \leq 1 \right\}$. If there exists a function $K \in L_1$,

$$K \sim \sum_{k=1}^{\infty} \lambda_k \cos \left(kx - \frac{\beta\pi}{2} \right),$$

Date: May 25, 2015.

2000 Mathematics Subject Classification. 42A10, 42A45.

Key words and phrases. Trigonometric system, n -width, optimal approximation.

then the set $\Lambda_\beta U_p$ can be represented in the convolution form

$$f(x) = \frac{1}{2\pi} \int_{\mathbb{T}^1} K(x-y) \phi(y) dy.$$

In this case we say that $f \in K * U_p$. If $K \notin L_1$ then we should consider generalised convolutions. In particular, if $\lambda_k = k^{-r}$, $\beta = r$, $r > 0$ we get Sobolev's classes W_p^r . If $\lambda_k = \exp(-\mu k^r)$, $\beta = 0$, $\mu > 0$, $0 < r < 1$, then the class $K * U_p$ consists of infinitely differentiable functions. In the case $\lambda_k = \exp(-\mu k^r)$, $\beta = 0$, $\mu > 0$, $r = 1$ ($r > 1$) we get classes of analytic (entire) functions respectively.

Let $\lambda_k = k^{-r}$, $\beta = r$. It is well-known that in this case ("finite smoothness", i.e. $r > 1/p$ if $1 < p \leq 2 \leq q < \infty$, or $2 \leq p \leq q < \infty$)

$$d_n(W_p^r, L_q) \asymp \begin{cases} n^{-r}, r > 0, & 1 < q < p < \infty, \\ n^{-r+1/p-1/q}, r > 1/p-1/q, & 1 < p \leq q \leq 2, \\ n^{-r+1/p-1/2}, r > 1/p, & 1 < p \leq 2 \leq q < \infty, \\ n^{-r}, r > 1/p, & 2 \leq p \leq q < \infty, \end{cases}$$

as $n \rightarrow \infty$. It is easy to show that

$$E_n(W_p^r, L_q) := \sup_{f \in W_p^r} \inf_{g \in \mathcal{T}_n} \|f - g\|_{L_q} \asymp n^{-r+(1/p+1/q)_+}, \quad r > 1/p-1/q,$$

where $(a)_+ := \max\{a, 0\}$. Hence the sequence of trigonometric polynomials $\mathcal{T}_n$ is optimal if $r > 1/p-1/q$, $1 < p \leq q \leq 2$, or $r > 0$, $1 < q < p < \infty$. In the remaining cases, $r > 1/p$, $1 < p \leq 2 \leq q < \infty$, $r > 1/p$, $2 \leq p \leq q < \infty$, $\mathcal{T}_n$ is not optimal. If $1/p-1/q < r < 1/p$ then we have the case of so-called "small smoothness",

$$d_n(W_p^r, L_q) \asymp n^\gamma, \quad 1/p-1/q < r < 1/p, \quad r \neq \beta, \quad 2 \leq p \leq q < \infty,$$

where

$$\gamma = \max\{-r, q(-r+1/p-1/q)/2\}$$

and

$$\beta = \frac{1/p-1/q}{2(1/2-1/q)}.$$

In the case

$$1/p-1/q < r < 1/p, \quad 1 < p < 2 < q < \infty$$

we have

$$d_n(W_p^r, L_q) \asymp n^{q(-r+1/p-1/q)/2}.$$

This means that the sequence of subspaces $\mathcal{T}_n$ is not optimal in this case. If $\lambda_k = \exp(-\mu k^r)$, $\beta = 0$, $\mu > 0$, $0 < r < 1$, then

$$d_{2n}(W_p^r, L_q) \asymp \begin{cases} \exp(-\mu n^r) n^{(1-r)(1/p-1/q)}, & 1 < p \leq q \leq 2, \\ \exp(-\mu n^r), & 1 < q \leq p \leq 2 \text{ or } 2 \leq p, q < \infty, \\ \exp(-\mu n^r) n^{(1-r)(1/p-1/2)} & 1 < p \leq 2 \leq q < \infty. \end{cases}$$

It is easy to show that

$$\begin{aligned} E_{2n}(K * U_p, L_q) &:= \sup_{f \in K * U_p} \inf_{g \in \mathcal{T}_n} \|f - g\|_{L_q} \\ &\asymp \exp(-\mu n^r) n^{(1-r)_+(1/p-1/q)}, \end{aligned}$$

$1 < p, q < \infty$, where

$$K(x) = \sum_{k=1}^{\infty} \exp(-\mu k^r) \cos kx.$$

Hence $\mathcal{T}_n$ is not optimal if $2 \leq p, q < \infty$ and $1 < p \leq 2 \leq q < \infty$.

Finally, in the case of analytic or entire smoothness, i.e. if $r \geq 1$ we have

$$d_{2n}(W_p^r, L_q) \asymp E_{2n}(K * U_p, L_q) \asymp \exp(-\mu n^r).$$

This means the the sequence of subspaces $\mathcal{T}_n$ is optimal for $K * U_p$ in L_q for any $1 < p, q < \infty$. See [5] for more information.

2. THE RESULTS

In this section we prove the following statement.

Theorem 1. *Let $\lambda_k = k^{-(1/p-1/q)+} (\ln k)^{-\gamma}$, $\gamma > 0$, $1 < p, q < \infty$, then the sequence of subspaces $\mathcal{T}_n$ is optimal in the sense of order as $n \rightarrow \infty$.*

Proof Let $\lambda_k = k^{-(1/p-1/q)+} (\ln k)^{-\gamma}$, $\gamma > 0$, $1 < p, q < \infty$. Using results from [4, 1] it is possible to show that

$$d_n(\Lambda U_p, L_q) \leq E_n(K * U_p, L_q) \ll (\ln k)^{-\gamma}, \quad 1 < p < q < \infty.$$

We turn to the lower bounds now. Clearly,

$$d_n(\Lambda U_p, L_q) \geq d_n(\Lambda U_p \cap \mathcal{T}_m, L_q)$$

and

$$d_n(\Lambda U_p \cap \mathcal{T}_m, L_q) \geq C_q^{-1} d_n(\Lambda U_p \cap \mathcal{T}_m, L_q \cap \mathcal{T}_m)$$

since $\|S_m : L_q \rightarrow L_q \cap \mathcal{T}_m\| \leq C_q$, $1 < q < \infty$, where S_m is the operator of orthogonal projection onto $\mathcal{T}_m$. Finally, applying Marcinkiewicz-Zygmund inequality

$$C_{1,p} \|t_m\|_p \leq \left(m^{-1} \sum_{k=1}^{2m+1} \left| t_n \left(\frac{2\pi k}{2m+1} \right) \right|^p \right)^{1/p} \leq C_{2,p} \|t_m\|_p, \quad \forall t_m \in \mathcal{T}_m$$

we get

$$d_n(\Lambda U_p, L_q) \gg (\ln m)^{-\gamma} d_n(B_p^m, l_q^m),$$

where the norm in l_q^m , $1 \leq q \leq \infty$ is defined as usual, $\|v\| := (\sum_{k=1}^m |v_k|^q)^{1/q}$ and B_p^m is the unit ball in l_p^m . We will need the following result [2, 3]

$$C_1(p, q) \leq \frac{d_n(B_p^m, l_q^m)}{\Phi(m, n, p, q)} \leq C_2(p, q)$$

for any $m > n$, where

$$\Phi(m, n, p, q) := \left(\min \left\{ 1, m^{1/q} n^{-1/2} \right\} \right)^{(1/p-1/q)/(1/2-1/q)},$$

if $2 \leq p < q \leq \infty$ and

$$\Phi(m, n, p, q) := \max \left\{ m^{1/q-1/p}, \min \left\{ 1, m^{1/q} n^{-1/2} \right\} (1 - n/m)^{1/2} \right\}$$

if $1 \leq p < 2 \leq q \leq \infty$. Let $2 \leq p < q \leq \infty$ then

$$\begin{aligned} d_n(\Lambda U_p, L_q) &\gg (\ln m)^{-\gamma} d_n(B_p^m, l_q^m) \\ &\gg (\ln m)^{-\gamma} \left(\min \left\{ 1, m^{1/q} n^{-1/2} \right\} \right)^{(1/p-1/q)/(1/2-1/q)}. \end{aligned}$$

Let $m = n^{q/2}$ then

$$d_n(\Lambda U_p, L_q) \gg (\ln m)^{-\gamma} \gg (\ln n)^{-\gamma}.$$

If $1 < p < 2 \leq q < \infty$ then

$$\begin{aligned} d_n(\Lambda U_p, L_q) &\gg (\ln m)^{-\gamma} \left(\min \left\{ 1, m^{1/q} n^{-1/2} \right\} \right)^{(1/p-1/q)/(1/2-1/q)} \\ &\gg (\ln m)^{-\gamma}. \end{aligned}$$

Hence the trigonometric system is an optimal in the logarithmic neighbourhood of embedding of L_p into L_q .

REFERENCES

- [1] Bordin, B., Kushpel, A., Levesley, J., Tozoni, S., Estimates of n -widths of Sobolev's classes on compact globally symmetric spaces of rank 1, *it Journ. of Func. Anal.* **202**, 2003, 307-326.
- [2] Kashin, B. S., n -Widths of some finite dimensional sets and classes of smooth functions, *Izvestiya Acad. Nauk. USSR*, **41**, 1977, 334-351.
- [3] Kashin, B. S., On n -widths of Sobolev's classes of small smoothness, *Vestnik MGU*, **5**, 1981, 50-54.
- [4] Kushpel, A. K., On an estimate of widths of sets in some function spaces, In: Some problems in the theory of approximation of functions, Kiev, Inst. Math. Acad. Nauk Ukrain. SSR, 1988, 92-95.
- [5] Kushpel, A. K., Optimal approximation on $\mathbb{S}^d$, *Journ. of Complexity*, **16**, 2000, 424-458.
- [6] Stepanets, A. I., Classification and approximation of periodic functions, Naukova Dumka, Kiev, 1987.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LEICESTER
E-mail address: `ak412@le.ac.uk`